

# Analytic Continuation of Two-Loop Hexa-Box Integrals

based on work with Costas Papadopoulos,  
in extension of [[arXiv:2201.07509](https://arxiv.org/abs/2201.07509) [[hep-ph](#)]]

**Nikos Dokmetzoglou**

National Centre for Scientific Research "Demokritos"  
Institute of Nuclear and Particle Physics

March 28, 2024

NCSR-D INPP Annual Meeting 2024



DEMOKRITOS

NATIONAL CENTRE FOR SCIENTIFIC RESEARCH



**H.F.R.I.**  
Hellenic Foundation for  
Research & Innovation

# Outline

- Hexa-Box Integral Families through the SDE Approach
- Mathematica Implementation
- Analytic Continuation

# Hexa-Box Integral Families through the SDE Approach

## Brief Review of the Simplified Differential Equations (SDE) Approach

In the SDE approach for Master Integrals (Papadopoulos 2014), the momenta are parametrized by introducing a dimensionless variable  $x$ , as follows

$$q_1 \rightarrow p_{123} - xp_{12}, \quad q_2 \rightarrow p_4, \quad q_3 \rightarrow -p_{1234}, \quad q_4 \rightarrow xp_1$$

where the new momenta  $p_i$ ,  $i = 1 \dots 5$  satisfy now  $\sum_1^5 p_i = 0$ ,  $p_i^2 = 0$ ,  $i = 1 \dots 5$ , whereas  $p_{i\dots j} := p_i + \dots + p_j$ . The set of independent invariants is given by  $\{S_{12}, S_{23}, S_{34}, S_{45}, S_{51}, x\}$ , with  $S_{ij} := (p_i + p_j)^2$ . The explicit mapping between the two sets of invariants is given by

$$q_1^2 = (1-x)(S_{45} - S_{12}x), \quad s_{12} = (S_{34} - S_{12}(1-x))x, \quad s_{23} = S_{45}, \quad s_{34} = S_{51}x, \\ s_{45} = S_{12}x^2, \quad s_{15} = S_{45} + (S_{23} - S_{45})x$$

and as usual the  $x = 1$  limit corresponds to the on-shell kinematics.

## Brief Review of the Simplified Differential Equations (SDE) Approach


For pure bases with rational alphabet letters or algebraic alphabet letters with rationalizable square roots, the differential equation for Master Integrals can be cast into its canonical form:

$$\partial_x \mathbf{g} = \epsilon \left( \sum_{i=1}^{l_{max}} \frac{\mathbf{M}_i}{x - l_i} \right) \mathbf{g} \quad (3)$$

However, for some topologies there are some alphabet letters with square roots that cannot be rationalized simultaneously, leading to the more general form of the SDE:

$$\partial_x \mathbf{g} = \epsilon \left( \sum_{a=1}^{l_{max}} \frac{d \log L_a}{dx} \mathbf{M}_a \right) \mathbf{g} \quad (4)$$

## 5-Point 2-Loop 1-Mass Non-Planar Topologies



**Figure:** The five non-planar families with one external massive leg. The first row corresponds to the so-called hexabox topologies, whereas the diagrams of the second row are known as double-pentagons. We label them as follows:  $N_1$  (top left),  $N_2$  (top middle),  $N_3$  (top right),  $N_4$  (bottom left),  $N_5$  (bottom right).

## Hexa-Box Topologies: Analytic up to Weight 2

For all hexa-box integral families, up to transcendental weight 1, all square roots appearing in the corresponding alphabet letters can be rationalized, and analytic expressions can thus be obtained up to weight 2, using the iterated integral definition of Goncharov Polylogarithms (GPLs):

$$\mathcal{G}(a_n, \dots, a_1; x) = \int_0^x dt \frac{1}{t - a_n} \mathcal{G}(a_{n-1}, \dots, a_1; t) \quad (5)$$

However, for topologies  $N_2$  and  $N_3$  non-rationalizable square roots break this procedure above weight 2.

## Hexa-Box Topologies: Semi-analytic above Weight 2

Above weight 2, we resort to a semi-numerical one-dimensional integral representation (Kardos, Papadopoulos, Smirnov, Syrrakos, Wever 2022). At weight 3,

$$\partial_x g_l^{(3)} = \sum_a (\partial_x \log L_a) \sum_J c_{IJ}^a g_J^{(2)} \quad (6)$$

$$g_l^{(3)} = g_{l,\mathcal{G}}^{(3)} + b_l^{(3)} + \int_0^{\bar{x}} dx \left( \sum_a (\partial_x \log L_a) \sum_J c_{IJ}^a g_J^{(2)} - \sum_a \frac{l_a}{x} \sum_J c_{IJ}^a g_{J,0}^{(2)} \right) \quad (7)$$

with  $b_l^{(3)}$  being the boundary terms at  $\mathcal{O}(\epsilon^3)$  and  $g_{l,\mathcal{G}}^{(3)} = \int_0^{\bar{x}} dx \sum_a \frac{l_a}{x} \sum_J c_{IJ}^a g_{J,0}^{(2)} \Big|_{\mathcal{G}}$ , with the subscript  $\mathcal{G}$ , indicating that the integral is represented in terms of GPLs.

## Hexa-Box Topologies: Semi-analytic above Weight 2

At weight 4,

$$\begin{aligned}
g_l^{(4)} = & g_{l,\mathcal{G}}^{(4)} + b_l^{(4)} + \left( \sum_a \log L_a \sum_J c_{IJ}^a g_J^{(3)} \right) - \left( \sum_a LL_a \sum_J c_{IJ}^a g_{J,0}^{(3)} \right) \\
& - \int_0^{\bar{x}} dx \sum_a (\log L_a - LL_a) \sum_J c_{IJ}^a \sum_b \frac{l_b}{x} \sum_K c_{JK}^b g_{K,0}^{(2)} \\
& - \int_0^{\bar{x}} dx \sum_a \log L_a \sum_J c_{IJ}^a \left( \sum_b (\partial_x \log L_b) \sum_K c_{JK}^b g_K^{(2)} - \sum_b \frac{l_b}{x} \sum_K c_{JK}^b g_{K,0}^{(2)} \right)
\end{aligned} \tag{8}$$

# Mathematica Implementation

## Mathematica Implementation

$$g_I^{(n+1)} = g_{I,\mathcal{G}}^{(n+1)} + b_I^{(n+1)} + \int_0^{\bar{x}} dx \left( \sum_a (\partial_x \log L_a) \sum_J c_{IJ}^a g_J^{(n)} - \sum_a \frac{l_a}{x} \sum_J c_{IJ}^a g_{J,0}^{(n)} \right)$$

## Mathematica Implementation

$$g_I^{(n+1)} = g_{I,\mathcal{G}}^{(n+1)} + b_I^{(n+1)} + \int_0^{\bar{x}} dx \left( \sum_a (\partial_x \log L_a) \sum_J c_{IJ}^a \boxed{g_J^{(n)}} - \sum_a \frac{l_a}{x} \sum_J c_{IJ}^a \boxed{g_{J,0}^{(n)}} \right)$$

**PureBasisElementAnalytic**[*ElementIndex*, *TranscendentalWeight*] : analytic form of pure basis elements, up to weight 2 for  $N_2$  and  $N_3$  hexa-box families

**PureBasisElementAnalytic**[*ElementIndex*, *TranscendentalWeight*,  $x \rightarrow 0$ ] :

$x \rightarrow 0$  limit of analytic form of pure basis elements, e.g.  $g_{J,0}^{(2)}$  are obtained by expanding  $g_J^{(2)}$  around  $x = 0$  and keeping terms up to order  $\mathcal{O}(\log(x)^2)$

## Mathematica Implementation

$$g_I^{(n+1)} = g_{I,\mathcal{G}}^{(n+1)} + b_I^{(n+1)} + \int_0^{\bar{x}} dx \left( \sum_a (\partial_x \log L_a) \sum_J c_{IJ}^a g_J^{(n)} - \sum_a \frac{l_a}{x} \sum_J c_{IJ}^a g_{J,0}^{(n)} \right)$$

**BoundaryTerm**[*ElementIndex*, *TranscendentalWeight*] : analytic form of boundary terms at  $\mathcal{O}(\epsilon^{(n+1)})$

# Mathematica Implementation

$$g_l^{(n+1)} = g_{l,\mathcal{G}}^{(n+1)} + b_l^{(n+1)} + \int_0^{\bar{x}} dx \left( \sum_a (\partial_x \log L_a) \sum_J c_{lJ}^a g_J^{(n)} - \sum_a \frac{l_a}{x} \sum_J c_{lJ}^a g_{J,0}^{(n)} \right)$$

**DLogAlphabetLetter**[*LetterIndex*] : derivatives of logs of alphabet letters with respect to parameter  $x$

**DLogAlphabetLetter**[*LetterIndex*,  $x \rightarrow 0$ ] :  $x \rightarrow 0$  limit of  $\partial_x \log L_a$ , obtained by expanding  $\partial_x \log L_a$  around  $x = 0$  and keeping terms up to order  $\mathcal{O}(x^{-1})$

**LogAlphabetLetter**[*LetterIndex*] : logs of alphabet letters

**LogAlphabetLetter**[*LetterIndex*,  $x \rightarrow 0$ ] :  $x \rightarrow 0$  limit of  $\log L_a$ , obtained by expanding  $\log L_a$  around  $x = 0$  and keeping terms up to order  $\mathcal{O}(\log(x))$

## Mathematica Implementation

$$g_l^{(n+1)} = g_{l,\mathcal{G}}^{(n+1)} + b_l^{(n+1)} + \int_0^{\bar{x}} dx \left( \sum_a (\partial_x \log L_a) \sum_J c_{lJ}^a g_J^{(n)} - \sum_a \frac{l_a}{x} \sum_J c_{lJ}^a g_{J,0}^{(n)} \right)$$

DifferentialEquationRHS[*ElementIndex*, *TranscendentalWeight*]

DifferentialEquationRHS[*ElementIndex*, *TranscendentalWeight*, x → 0]

## Mathematica Implementation

$$g_l^{(n+1)} = \boxed{g_{l,\mathcal{G}}^{(n+1)}} + b_l^{(n+1)} + \int_0^{\bar{x}} dx \left( \sum_a (\partial_x \log L_a) \sum_J c_{IJ}^a g_J^{(n)} - \sum_a \frac{l_a}{x} \sum_J c_{IJ}^a g_{J,0}^{(n)} \right)$$

$$\boxed{g_{l,\mathcal{G}}^{(n+1)} = \int_0^{\bar{x}} dx \sum_a \frac{l_a}{x} \sum_J c_{IJ}^a g_{J,0}^{(n)} \Big|_{\mathcal{G}}}$$

**ClassicalToGoncharovPolyLogs[*expr*]** and **GoncharovToClassicalPolyLogs[*expr*]** :  
converter between Goncharov Polylogarithms and Classical Logs and PolyLogs, making  
use of ([Frellesvig et al. 2016](#))

Analytic integration performed using the function **GIntegrate** of the PolyLogTools  
package ([Duhr, Dulat 2019](#)).

## Mathematica Implementation

$$g_I^{(n+1)} = g_{I,G}^{(n+1)} + b_I^{(n+1)} + \int_0^{\bar{x}} dx \left( \sum_a (\partial_x \log L_a) \sum_J c_{IJ}^a g_J^{(n)} - \sum_a \frac{l_a}{x} \sum_J c_{IJ}^a g_{J,0}^{(n)} \right)$$

**PureBasisElementNumerical**[*ElementIndex*, *TranscendentalWeight*, *PhaseSpacePoint*] : numerical evaluation of pure basis elements at weight  $n + 1$  by integration of the analytic form of pure basis elements at weight  $n$

# Mathematica Implementation

Other auxiliary functions:

- `LettersToInvariants[expr]` and `InvariantsToLetters[expr]`
- `AssignImaginaryParts[expr, PhaseSpacePoint]`
- `NumericalEvaluationMathematica[expr, PhaseSpacePoint]` and `NumericalEvaluationGiNaC[expr, PhaseSpacePoint]`

# Analytic Continuation

## Need for Analytic Continuation

- For an arbitrary kinematic configuration  $(S_{12}, S_{23}, S_{34}, S_{45}, S_{51})$ , possible singularities in the domain of integration  $(0, \bar{x})$
- Need to analytically continue the integrand of the numerical integration, i.e. the differential equation at weight  $n + 1$  and its  $x \rightarrow 0$  limit
- Types of singularities encountered:
  - $g_J^{(n \leq 2)}$  and  $g_{J,0}^{(n \leq 2)}$  : logarithmic and polylogarithmic branch points/cuts
  - $\partial_x \log L_a$  : poles at points  $x = \ell_i$
  - $\log L_a$  and  $LL_a$  (the  $x \rightarrow 0$  limits of  $\log L_a$ ) : logarithmic branch points/cuts
  - square-root branch points/cuts

## Imaginary Parts from $\mathcal{F}$ Symanzik Polynomial

The ordinary momentum-space representation of Feynman integrals:

$$G(\nu_1, \dots, \nu_n) = \int \prod_{l=1}^L \frac{d^d k_l}{i\pi^{d/2}} \prod_{j=1}^n \frac{1}{D_j^{\nu_j}(\{k\}, \{p\}, m_j^2)}$$

The Feynman parameter representation of Feynman integrals:

$$G(\nu_1, \dots, \nu_n) = \Gamma\left(\nu - \frac{Ld}{2}\right) \prod_{j=1}^n \left[ \int_0^\infty dx_j \frac{x_j^{\nu_j-1}}{\Gamma(\nu_j)} \right] \delta\left(1 - \sum_{j=1}^n x_j\right) \frac{\mathcal{U}^{\nu-(L+1)d/2}}{\mathcal{F}^{\nu-Ld/2}}$$

where the polynomials  $\mathcal{U}, \mathcal{F}$  are known as first and second Symanzik polynomials.

## Imaginary Parts from $\mathcal{F}$ Symanzik Polynomial

Since the  $\mathcal{F}$  Symanzik polynomial maintains the sign of the  $i0$  prescription of Feynman propagators with all original invariants ( $s_{ij}$ ), assuming  $s_{ij}(p_{1s}) \rightarrow s_{ij}(p_{1s}) + i\eta$ , we determine the corresponding infinitesimal imaginary parts for our invariants ( $S_{ij}$ ) by recasting the  $\mathcal{F}$  Symanzik polynomial for a given integral family in terms of our invariants, extending those by  $S_{ij} \rightarrow S_{ij} + i\delta_{ij}\eta$ , and imposing a positivity constraint on the coefficients of all the Feynman parameters. For example, the positivity constraints for the  $N_2$  hexa-box integral family are:

$$\begin{aligned}\delta_{45} + \delta_{23}x &\geq \delta_{45}x, \delta_{51}x \geq 0, (x-1)(\delta_{12}x - \delta_{45}) \geq 0, \\ x(\delta_{34} + \delta_{12}(x-1)) &\geq 0, \delta_{12}x^2 \geq 0, \delta_{45} \geq 0, \\ x(\delta_{12} - \delta_{34} + \delta_{51}) &\leq 0, \delta_{23}x \geq x(\delta_{45} + \delta_{51})\end{aligned}$$

## Analytic Continuation of $\partial_x \log L_a$

As mentioned earlier,  $\partial_x \log L_a$  have poles at points  $x = \ell_j$ . To control the numerical integration over the locations of these poles, we make this pole structure manifest

$$\frac{d \log L_a}{dx} \rightarrow \frac{f(x)}{\prod_i (x - \ell_i)},$$

where  $f(x) \in \mathbb{R}$ , except for factors of  $\sqrt{\Delta} = i\sqrt{|\Delta|}$  for  $\Delta < 0$ , and we assign imaginary parts to all  $\ell_i$ 's using those we assigned to the  $S_{ij}$  invariants

$$\ell_i(S_{ij}) \rightarrow \ell_i(S_{ij} + i \delta_{ij} \eta) \equiv \ell_i + i \delta_i \eta.$$

## Analytic Continuation of $g_j^{(n \leq 2)}$

Concerning  $g_j^{(n \leq 2)}$ , their logarithmic and polylogarithmic branch points/cuts appear through expressions of the form:

- $\log(x)$ ,  $\log(l_a)$ ,  $\log(1 - l_a)$  or  $\log(l_3 - l_a)$ ,
- $\mathcal{G}(l_a; x)$ ,  $\mathcal{G}(0; x)$ ,  $\mathcal{G}(l_a, l_b; x)$ ,  $\mathcal{G}(0, l_a; x)$ ,  $\mathcal{G}(0, 0; x)$ ,

In elements  $g_{11}^{(2)}$  and  $g_{73}^{(2)}$  of the  $N_2$  hexa-box family we also find:

- $\mathcal{G}(0, 1; \tilde{l}_a(x))$  or  $\mathcal{G}(0, l_3; \tilde{l}_a(x))$ , and  $\mathcal{G}(1; \tilde{l}_a(x))$  or  $\mathcal{G}(l_3; \tilde{l}_a(x))$ ,

where  $\tilde{l}_a(x)$  are **algebraic expressions of  $x$** . To control the numerical integration over the locations of all branch points, we assign imaginary parts to all  $l_i$ 's similarly to before, and also to the  $\tilde{l}_a(x)$ 's, such that

$$\tilde{l}_a(l_i) \rightarrow \tilde{l}_a(l_i + i \delta_i \eta) \equiv \tilde{l}_a(l_i) + i \delta_a \eta.$$

Analytic Continuation of  $g_J^{(n \leq 2)}$ 

$$\tilde{\ell}_a(x) \in \left\{ \frac{x(S_{12}x - S_{23} - S_{34} + 2S_{51}) + S_{45} \pm \sqrt{\Delta(1, x)}}{2x(S_{12} - S_{34} + S_{51})}, \right. \\ \left. \frac{S_{12}S_{51}x^2 + x(-S_{12}S_{23} + 2S_{12}S_{45} + S_{23}S_{34} - S_{34}S_{45}) + S_{45}S_{51} \pm \sqrt{\Delta(2, x)}}{2S_{45}(S_{12} - S_{34} + S_{51})} \right\}$$

## Analytic Continuation of $\log L_a$

As for the logs of the alphabet letters, we isolate their logarithmic branch points in the following way:

$$\log L_a \rightarrow \log \left( L_a \frac{\prod_{i_D} (x - \ell_{i_D})}{\prod_{j_N} (x - \ell_{j_N})} \right) - \log \left( \prod_{i_D} (x - \ell_{i_D}) \right) + \log \left( \prod_{j_N} (x - \ell_{j_N}) \right)$$

where  $\left( L_a \frac{\prod_{i_D} (x - \ell_{i_D})}{\prod_{j_N} (x - \ell_{j_N})} \right) \in \mathbb{R}$ , except for factors of  $\sqrt{\Delta} = i\sqrt{|\Delta|}$  for  $\Delta < 0$ , and has no zeroes or poles in  $x \in (0, \bar{x})$ , and then we assign imaginary parts in the ordinary way to all  $\ell_j$ 's.

## Summary and Outlook

- Need for semi-analytic approach above transcendental weight 2 for some of the non-planar 5-point topologies
- Need for analytic continuation of the integrands
- Mathematica implementation, successful check for  $N_2$  integrals family on a specific physical phase-space point, against the numerical results obtained from the literature ([Abreu, Ita, Page, Tschernow 2022](#)), up to weight 3
- Need to extend our Mathematica implementation to weight 4, as well as to the other non-planar topologies
- More work to be done on generalizing our analytic continuation methods

# Thank you for your attention!

This research is supported by the Hellenic Foundation for Research and Innovation (H.F.R.I.), under the 2nd Call for H.F.R.I. Research Projects for the Support of Faculty Members and Researchers (Project Acronym: HOCTools-II, Project Number: 2674).



# References

- Abreu, S., Ita, H., Page, B., Tschernow, W. (2022). Two-loop hexa-box integrals for non-planar five-point one-mass processes, *JHEP* **03**: 182.
- Duhr, C., Dulat, F. (2019). PolyLogTools - Polylogs for the masses, *JHEP* **08**(8): 135.
- Frellesvig, H., Tommasini, D., Wever, C. (2016). On the reduction of generalized polylogarithms to  $\text{Li}_n$  and  $\text{Li}_{\{2,2\}}$  and on the evaluation thereof, *JHEP* **03**: 189.
- Kardos, A., Papadopoulos, C. G., Smirnov, A. V., Syrrakos, N., Wever, C. (2022). Two-loop non-planar hexa-box integrals with one massive leg, *JHEP* **05**(5): 033.
- Papadopoulos, C. G. (2014). Simplified differential equations approach for Master Integrals, *JHEP* **07**: 088.